

# Vibrazioni a onde meccanosonore

Paolo Buselli<sup>1</sup>, S. Messina<sup>2</sup>, M. Fontanesi<sup>3</sup>, F. Ferraro<sup>4</sup>, V. Rosati<sup>5</sup>,  
Maria Grazia Benedetti<sup>6</sup>, S. Filoni<sup>7</sup>, Roberto Casale<sup>8</sup>



## BIBLIOGRAFIA

- HAGBARTH KE, EKLUND G. Tonic vibration reflexes (TVR) in spasticity. *Brain Res* 1966; 2: 201-03.
- HAGBARTH KE, EKLUND G. The effects of muscle vibration in spasticity, rigidity, and cerebellar disorders. *J Neurol Neurosurg Psychiat* 1968; 31: 207-13.
- JOHNSTON RM, BISHOP B, COFFEY GH. Mechanical vibration of skeletal muscles. *Phys Ther* 1970; 50(4): 499-505.
- ARCANGEL CS, JOHNSTON R, BISHOP B. The achilles tendon reflex and the H response during and after tendon vibration. *Phys Ther* 1971; 51(8): 899-905.
- ARMSTRONG TJ, FINE LJ, RADWIN RG, SILVERSTEIN BS. Ergonomics and the effects of vibration in hand-intensive work. *Scand J Work Environ Health* 1987; 13(4): 286-9.
- MATYAS TA, GALEA MP, SPICER SD. Facilitation of the maximum voluntary contraction in hemiplegia by concomitant cutaneous stimulation. *Am J Phys Med* 1986; 65(3): 125-34.
- SAMUELSON B, JORFELDT L, AHLBORG B. Influence of vibration on endurance of maximal isometric contraction. *Clin Physiol* 1989; 9(1): 21-5.
- CASALE R, RING H, RAINOLDI A. High frequency vibration conditioning stimulation centrally reduces myoelectrical manifestation of fatigue in healthy subjects. *J Electromyogr Kinesiol* 2009; 19: 998-1004.
- PIETRANGELO T, MANCINELLI R, TONIOLO L, ET AL. Effects of local vibrations on skeletal muscle trophism in elderly people: mechanical, cellular, and molecular events. *Int J Molecular Med* 2009; 24: 503-12.
- TSUJI T, KITANO N, TSUNODA K, ET AL. Short-term effects of whole-body vibration on functional mobility and flexibility in healthy, older adults: a randomized crossover study. *J Geriatr Phys Ther* 2014; 37(2): 58-64.
- CARDINALE M, BOSCO C. The use of vibration as an exercise intervention. *Exerc Sport Sci Rev* 2003; 31(1): 3-7.
- MARÍN PJ, RHEA MR. Effects of vibration training on muscle power: a metaanalysis. *J Strength Cond Res* 2010; 24(3): 871-8.
- NOMA T, MATSUMOTO S, ETOH S., Shimodozono M., Kawahira K. Anti-spastic effects of the direct application of vibratory stimuli to the spastic muscles of hemiplegic limbs in post stroke patients. *Brain Inj* 2009; 23 (7-8): 623-31.
- NOMA T, MATSUMOTO S, ETOH S, SHIMODOZONO M, KAWAHIRA K. Anti-spastic effects of the direct application of vibratory stimuli to the spastic muscles of hemiplegic limbs in post-stroke patients: a proof-of-principle study. *J Rehabil. Med* 2012; 44: 325-30.
- PANG MYC, LAU RWK, YIP SP. The effects of whole-body vibration therapy on bone turnover, muscle strength, motor function, and spasticity in chronic stroke: a randomized controlled trial. *Eur J Phys Rehabil Med.* 2013; 49: 1-12.
- ROSENKRANZ K, PESENTI A, PAULUS W, TERGAU F. Focal reduction of intracortical inhibition in the motorcortex by selective proprioceptive stimulation. *Exp Brain Res* 2003; 149(1): 9-16.
- FORNER-CORDERO A, STEYVERS M, LEVIN O, ALBERTS K, SWINNEN SP. Changes in corticomotor excitability following prolonged muscle tendon vibration. *Behav Brain Res* 2008; 190(1): 41-9.
- MARCONI B, FILIPPI GM, KOCH G, ET AL. Long-term effects on cortical excitability and motor recovery induced by repente muscle vibration in chronic stroke patients. *Neurorehabil Neural Repair* 2010; 25(1): 48-60.
- CASALE R, DAMIANI C, MAESTRI R, ET AL. Localized 100Hz vibration improves function and reduces upper limb spasticity: a double-blind controlled study. *Eur J Phys Rehabil Med* 2014 Mar 21 [Epub ahead of print].
- LIEPERT J, BINDER C. Vibration-induced effects in stroke patients with spastic hemiparesis – a pilot study. *Resto Neurol Neurosci* 2010; 28: 729-35.
- CONRAD MO, SCHEIDT RA, SCHMIT BD. Effects of wrist tendon vibration on arm tracking in people post stroke. *J Neurophysiol* 2011; 106: 1480-8.
- CALIANDRO P, CELLETTI C, PADUA L, ET AL. Focal muscle vibration in the treatment of upper limb spasticity: a pilot randomized controlled trial in patients with chronic stroke. *Arch Phys Med Rehabil* 2012; 93: 1656-61.
- TAVERNESE E, PAOLONI M, MANGONE M, ET AL. Segmental muscle vibration improves reaching movement in patients with chronic stroke. A randomized controlled trial. *NeuroRehabilitation* 2013; 32(3): 591-9.
- PAOLONI M, GIOVANNELLI M, MANGONE M, ET AL. Does living segmental muscle vibration alter the response to botulinum toxin injections in the treatment of spasticity in people with multiple sclerosis? A single-blind randomized controlled trial. *Clin Rehabil* 2013; 27(9): 803-12.

25. CASALE R, TIENGO M. Flexion withdrawal reflex: a link between pain and Motility. In: M. Tiengo et al. (Eds) *Advance in pain research and therapy*. New York: Raven Press, , 1987; 10: 77-83.
26. KAKIGI R, SHIBASAKI H. Mechanisms of pain relief by vibration and movement. *J Neurol Neurosurg Psychiatry* 1992; 55(4): 282-6.
27. ANNINO G, PADUA E, CASTAGNA C, ET AL. Effect of whole body vibration training on lower limb performance in selected high-level ballet students. *J Strength Cond Res* 2007; 21(4): 1072-6.
28. MARÍN PJ, HAZELL TJ, GARCÍA-GUTIÉRREZ MT, COCHRANE DJ. Acute unilateral leg vibration exercise improves contralateral neuromuscular performance. *J Musculoskelet Neuronal Interact* 2014; 14(1): 58-67.
29. DAVIS BL, CAVANAGH PR, PERRY JE. Locomotion in a rotating space station: a synthesis of new data with established concepts. *Gait Posture* 1994; 2: 157-65.
30. PAOLONI M, MANGONE M, SCETTRI P, ET AL. Segmental muscle vibration improved walking in chronic stroke patients with foot drop: a randomized controlled trial. *Neurorehabil Neural Repair* 2010; 24(3): 254-62. ■